

The Open University of Sri Lanka  
B.Sc. / B.Ed. Degree Programme  
Final Examination -2019/2020  
Applied Mathematics-- Level 05  
ADU5307– Numerical Methods



Duration: Two Hours

Date: 27. 12. 2019

Time: 02.00 p.m. – 4.00 p.m.

Answer Four Questions Only.

1. (a) Show that recurrence relation to find the root of a given equation by method of False

Position is given by  $x_{n+1} = x_{n-1} - \frac{f(x_{n-1})}{f(x_n) - f(x_{n-1})}(x_n - x_{n-1})$  where  $n = 1, 2, 3, \dots$

- (b) Show that the equation  $x^4 + x^2 - 80 = 0$ , has a root in the interval  $[2, 3]$  and use method of False Position to find the root correct to four decimal places.

2. (a) Prove that

(i)  $E = (1 - \nabla)^{-1}$ ,

(ii)  $\delta = E^{1/2} - E^{-1/2}$ ,

(iii)  $\mu = \frac{1}{2}(E^{1/2} + E^{-1/2})$  where  $\nabla$ ,  $\delta$ ,  $E$  and  $\mu$  are the backward difference, the central difference, the shift and the average operators respectively.

- (b) Derive Gregory- Newton backward interpolation formula.

- (c) The table given below shows the melting point of an alloy of lead and zinc, where  $y$  is the temperature in  $^{\circ}\text{C}$  and  $x$  is the percentage of lead in the alloy.

$x\%$	40	50	60	70	80	90
$y\ ^{\circ}\text{C}$ ,	184	204	226	250	276	304

Using suitable interpolation formula, find the melting point of the alloy containing 84% of lead.

3. (a) Derive Cubic Spline interpolation formula

$$M_{i-1} + 4M_i + M_{i+1} = \frac{6}{h^2}(y_{i-1} - 2y_i + y_{i+1}), \text{ for } i = 1 \text{ to } n-1, \text{ where } f''(x_i) = M_i, \text{ and}$$

$$x_{i+1} - x_i = h.$$

(b) Following values of  $x$  and  $y$  are given.

$x$	1	2	3	4
$y$	1	2	5	11

Using Cubic Spline interpolation method evaluate  $y(1.5)$ .

4. (a) Derive the Simpson's One-Third Rule.

(b) If the interval  $[a, b]$  is divided into  $2n$  sub intervals and corresponding ordinates are denoted by  $y_0, y_2, \dots, y_{2n}$  then show that the error in Simpson's One-Third rule is given by  $|E| < \frac{(b-a)h^4}{180} M$ , where  $M$  is the numerically greater value of  $y_0^{iv}, y_2^{iv}, \dots, y_{2n-2}^{iv}$ .

(c) Evaluate  $\int_0^1 \frac{x^2}{1+x^3} dx$  using Simpson's One-Third rule with  $h = 0.25$ . Hence find an approximate value for  $\ln 2^{1/3}$ .

5. (a) (i) Derive formula for Picard's method to solve  $\frac{dy}{dx} = f(x, y)$  subject to the initial condition

$$y(x_0) = y_0.$$

(ii) Using Picard's method, find the first-three successive approximations to solve

$$\frac{dy}{dx} = 2x - y^2 \text{ with the initial condition } y(0) = 0.$$

(b) Applying Euler's method with step size  $h = 0.1$ , find the value of  $x(0.5)$  for the initial value problem  $\frac{dy}{dx} = \frac{4-x}{x+y}$  subject to the initial condition  $y(0) = 1$ .

6. (a) State fourth order Runge-Kutta algorithm to solve  $\frac{dy}{dx} = f(x, y)$  subject to the initial condition  $y(x_0) = y_0$ .
- (b) Using Runge-Kutta method of fourth order, solve  $\frac{dy}{dx} = x + y$  subject to the initial condition  $y(0) = 1$ , at  $x = 0.1, 0.2$  and  $0.3$ .
- (c) Solve the second order differential equation  $\frac{d^2y}{dx^2} = y^3$  with the initial condition  $y(0) = 10$ ,  $y'(0) = 5$  using Runge-Kutta method of fourth order and evaluate  $y(0.1)$ .

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18